

EXISTENCE OF SOLUTION OF INTEGRAL EQUATIONS IN CONE METRIC SPACES

Shiv Kant Tiwari

Department of Mathematics, Lukhdhirji Engineering College, Morbi (Gujarat)-363642, India

Email: shivkant.math@gmail.com

Received on: 03/09/2022

Accepted on: 26/07/2023

Abstract

In this study, we use comparison mapping for contractive conditions in cone metric spaces to demonstrate the existence of a solution to mixed type integral equations. We use examples to show how our findings can be put to use.

Keywords: Integral equation, Cone metric space, Contractive mapping.

MSC (2020): 47H10, 54E50, 47G20, 34K20

1. Introduction

Fractional The French mathematician Frechet [15] originally described and investigated an abstract metric space in 1906. Numerous researchers [6, 8, 10, 18, 19, 20] have expanded the definition of metric space to include cone, semi, and quasi metric spaces, as well as generalized contraction mappings with applications.

One of the most vital areas of mathematics is fixed point theory, with numerous applications in several aspects of physical science as well as various domains of mathematics. The Banach contraction principle has a large number of generalizations like Ciric contraction, S-contraction, Chatterjee contraction, weak contractions principle etc. These contractions proved many results in analysis and play a key part in determining the existence and uniqueness of the findings in analysis and many other areas of mathematics.

Branciari [3] derived results in a fixed point theory for a single mapping in 2002. The authors [1, 2, 9, 11, 12, 16] used integral type equations to demonstrate several fixed point theorems.

2. Preliminaries

Definition 2.1 [7, 13, 14] Assume that B is a real Banach space and consider a subset $\mathbb{C}$ of B is referred to as a cone if satisfy:

1. $\mathbb{C} \neq \{0\}$, nonempty and closed
2. $a, b \in \mathbb{R}^+, \alpha, \beta \in \mathbb{C} \Rightarrow a\alpha + b\beta \in \mathbb{C}$;
3. $\alpha \in \mathbb{C}$ and $-\alpha \in \mathbb{C} \Rightarrow \alpha = 0$.

A partial order relation $\leq$ on a cone $\mathbb{C}$ define as $\alpha \leq \beta$ if and only if $\beta - \alpha \in \mathbb{C}$. If there is a positive number $a > 0$ such that

$$0 \leq \alpha \leq \beta \Rightarrow \|\alpha\| \leq a \|\beta\|,$$

for all $\alpha, \beta \in B$, then cone $\mathbb{C}$ is known as normal and the least positive value which holds the above inequality is known as normal constant of cone $\mathbb{C}$.

Definition 2.2 [7, 13, 14] Consider B be a Banach space and X be a nonempty subset of B , then a mapping $d: X \times X \rightarrow B$ is known as cone and (X, d) is known as cone metric space if satisfies:

- (i) $d(\alpha, \beta) \geq 0$ and $d(\alpha, \beta) = 0$ if and only if $\alpha = \beta$;
- (ii) $d(\alpha, \beta) = d(\beta, \alpha)$;
- (iii) $d(\alpha, \beta) \leq d(\alpha, \gamma) + d(\gamma, \beta)$

for all $\alpha, \beta, \gamma \in X$.

Example 2.1 [13, 17] Consider the supremum norm on the the Banach space $B = \mathbb{C}([0, l], \mathbb{Z})$, which are all the continuous functions from $[0, l]$ into $\mathbb{Z}$, such that

$$\|\alpha\|_{\infty} = \sup\{\|\alpha(s)\| : s \in [0, l]\}.$$

Assume $\mathbb{C} = \{(\alpha, \beta) : \alpha, \beta \geq 0\} \subset B = \mathbb{R}^2$ and $d(\zeta, \xi) = (\|\zeta - \xi\|, \kappa \|\zeta - \xi\|_{\infty})$, for every $\zeta, \xi \in B$. Then (B, d) is a cone metric space on the supremum norm.

Definition 2.3 [17] Consider a function $\Psi: X \rightarrow X$ on an ordered space X is said to be comparison function if it implies $\alpha \leq \beta$, $\Psi(\alpha) \leq \Psi(\beta)$, $\Psi(\alpha) \leq \alpha$ and $\lim_{n \rightarrow \infty} \|\Psi^n(\alpha)\| = 0$, for every $\alpha, \beta \in X$.

Example 2.2 [17] Let $\mathbb{C} = \{(\alpha, \beta) \in B: \alpha, \beta \geq 0\}$, where $B = \mathbb{R}^2$, then a mapping $\Psi: B \rightarrow B$ defined as $\Psi(\alpha, \beta) = (a\alpha, a\beta)$ is a comparison function for $a \in (0, 1)$. If Ψ_1, Ψ_2 are two comparison functions over $\mathbb{R}$, then $\Psi(\alpha, \beta) = (\Psi_1(\alpha), \Psi_2(\beta))$ is also a comparison function over B .

Lemma 2.1 [17] Given that normal constant β on a normal cone $\mathbb{C}$. Let $f: X \rightarrow X$ be a function on a complete cone metric space (X, d) as follows

$$d(f(\alpha), f(\beta)) \leq \Psi(d(\alpha, \beta)),$$

for all $\alpha, \beta \in X$. If it exists as a comparison function $\Psi: \mathbb{C} \rightarrow \mathbb{C}$, then f has a unique fixed point.

The aim of the current research is to generalize the findings of Pachpatte [4, 5] and numerous other articles by investigating the existence and uniqueness of solutions to the integrodifferential equations using fixed point theory on the cone metric spaces.

3. Main Results

Theorem 3.1 Assume the integrodifferential equations on a complete cone metric space (X, d) as follows

$$u(s) = f(s) + \int_0^s \zeta(s, t, u(t))dt + \int_0^l \xi(s, t, u(t))dt, \quad s \in [0, l], \quad (3.1)$$

$$u'(s) = f(s) + \int_0^s \zeta(s, t, u(t))dt + \int_0^l \xi(s, t, u(t))dt, \quad u(0) = u_0, \quad (3.2)$$

where $f: [0, l] \rightarrow \mathbb{Z}$, $\zeta, \xi: [0, l] \times [0, l] \times \mathbb{Z} \rightarrow \mathbb{Z}$ are continuous functions and $\mathbb{Z}$ is a Banach space on the norm $\|\cdot\|$, provided $u_0 \in \mathbb{Z}$. If the following criteria are met by integrodifferential equations:

($\mathbb{C}_1$) A comparison function $\Psi: \mathbb{R}^+ \rightarrow \mathbb{R}^+$ and continuous functions $q_1, q_2: [0, l] \times [0, l] \rightarrow \mathbb{R}^+$ are exists, such that

$$(\|\zeta(s, t, p) - \zeta(s, t, q)\|, \beta \|\zeta(s, t, p) - \zeta(s, t, q)\|) \leq q_1(s, t)\Psi(d(p, q)),$$

$$(\|\xi(s, t, p) - \xi(s, t, q)\|, \beta \|\xi(s, t, p) - \xi(s, t, q)\|) \leq q_2(s, t)\Psi(d(p, q)),$$

for every $s, t \in [0, l]$ and $p, q \in \mathbb{Z}$.

$$(\mathbb{C}_2) \quad \sup_{s \in [0, l]} \int_0^l [q_1(s, t) + q_2(s, t)] dt = 1, \int_0^l \int_0^l [q_1(s, t) + q_2(s, t)] dt \, ds \leq 1.$$

Then the unique solution of integrodifferential equations (3.1) and (3.2) are exists on $[0, l]$.

Proof : Consider the operator $\Gamma: B \rightarrow B$ is

$$\Gamma u(s) = f(s) + \int_0^s \zeta(s, t, u(t)) dt + \int_0^l \xi(s, t, u(t)) dt, \quad s \in [0, l]. \quad (3.3)$$

By conditions $\mathbb{C}_1$ and $\mathbb{C}_2$, we have

$$\begin{aligned} & (\| \Gamma u(s) - \Gamma v(s) \|, \beta \| \Gamma u(s) - \Gamma v(s) \|) \\ \leq & (\| \int_0^s \zeta(s, t, u(t)) dt + \int_0^l \xi(s, t, u(t)) dt - \int_0^s \zeta(s, t, v(t)) dt - \int_0^l \xi(s, t, v(t)) dt \|, \\ & \beta \| \int_0^s \zeta(s, t, u(t)) dt + \int_0^l \xi(s, t, u(t)) dt - \int_0^s \zeta(s, t, v(t)) dt - \int_0^l \xi(s, t, v(t)) dt \|) \\ \leq & (\int_0^s \| \zeta(s, t, u(t)) - \zeta(s, t, v(t)) \| dt + \int_0^l \| \xi(s, t, u(t)) - \xi(s, t, v(t)) \| dt, \\ & \beta \int_0^s \| \zeta(s, t, u(t)) - \zeta(s, t, v(t)) \| dt + \beta \int_0^l \| \xi(s, t, u(t)) - \xi(s, t, v(t)) \| dt) \\ \leq & (\int_0^s \| \zeta(s, t, u(t)) - \zeta(s, t, v(t)) \| dt, \beta \int_0^s \| \zeta(s, t, u(t)) - \zeta(s, t, v(t)) \| dt) \\ & + (\int_0^l \| \xi(s, t, u(t)) - \xi(s, t, v(t)) \| dt, \beta \int_0^l \| \xi(s, t, u(t)) - \xi(s, t, v(t)) \| dt) \\ \leq & \int_0^s q_1(s, t) \Psi(\| u - v \|_\infty, \beta \| u - v \|_\infty) dt + \int_0^l q_2(s, t) \Psi(\| u - v \|_\infty, \beta \| u - v \|_\infty) dt \\ \leq & \int_0^l q_1(s, t) \Psi(\| u - v \|_\infty, \beta \| u - v \|_\infty) dt + \int_0^l q_2(s, t) \Psi(\| u - v \|_\infty, \alpha \| u - v \|_\infty) dt \\ \leq & \int_0^l [q_1(s, t) + q_2(s, t)] \Psi(\| u - v \|_\infty, \beta \| u - v \|_\infty) dt \\ \leq & \Psi(\| u - v \|_\infty, \beta \| u - v \|_\infty) \int_0^l [q_1(s, t) + q_2(s, t)] dt \\ = & \Psi(\| u - v \|_\infty, \beta \| u - v \|_\infty), \end{aligned}$$

This means that

$$d(\Gamma u, \Gamma v) \leq \Psi(d(u, v))$$

for every $u, v \in B$. Now, the Lemma 2.1 has been applied, therefore the operator Γ has a unique fixed point in B and hence a unique solution of equations (3.1) and (3.2). Thus Theorem 3.1 has proved.

Example 3.1 Consider the mappings ζ and ξ such that

$$\zeta(s, t, u) = st + \frac{ut}{2}, \quad h(s, t, u) = (st)^2 + \frac{stu^2}{2}, \quad s, t \in [0, 1] \quad u \in \mathbb{C}([0, 1], \mathbb{R}), \quad (3.4)$$

and a metric $d(u, v)$ described as

$$d(u, v) = (\|u - v\|_\infty, \alpha \|u - v\|_\infty)$$

on $\mathbb{C}([0, 1], \mathbb{R})$ and $\beta \geq 0$. Then $\mathbb{C}([0, 1], \mathbb{R})$ is a complete cone metric space. We choose two continuous functions $q_1^*(s, t), q_2^*(s, t): [0, 1] \times [0, 1] \rightarrow \mathbb{R}^+$, such that $q_1^*(s, t) = t$ and $q_2^*(s, t) = st$ and a comparison function $\Psi^*: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ such that $\Psi^*(u, v) = \frac{1}{2}(u, v)$. Now, we demonstrate that

$$\begin{aligned} & (|\zeta(s, t, u(t)) - \zeta(s, t, v(t))|, \beta |\zeta(s, t, u(t)) - \zeta(s, t, v(t))|) \\ &= (|st + \frac{ut}{2} - st - \frac{vt}{2}|, \beta |st + \frac{ut}{2} - st - \frac{vt}{2}|) \\ &= (|\frac{ut}{2} - \frac{vt}{2}|, \beta |\frac{ut}{2} - \frac{vt}{2}|) \\ &= \frac{t}{2}(|u - v|, \beta |u - v|) \\ &\leq \frac{t}{2}(\|u - v\|_\infty, \beta \|u - v\|_\infty) \\ &= q_1^* \Psi^*(\|u - v\|_\infty, \beta \|u - v\|_\infty), \end{aligned}$$

Similarly, we can show that

$$(|\xi(s, t, u(t)) - \xi(s, t, v(t))|, \beta |\xi(s, t, u(t)) - \xi(s, t, v(t))|) \leq q_2^* \Psi^*(\|u - v\|_\infty, \beta \|u - v\|_\infty),$$

Now, we obtain

$$\begin{aligned} \int_0^1 [q_1^*(s, t) + q_2^*(s, t)] dt &= \int_0^1 [t + st] dt = \frac{1}{2}(1 + s) \Rightarrow \sup_{s \in [0, 1]} \left\{ \frac{1}{2}(1 + s) \right\} = 1. \\ \int_0^1 \int_0^1 [q_1^*(s, t) + q_2^*(s, t)] dt ds &= \int_0^1 \int_0^1 [t + st] dt ds = \int_0^1 \frac{1}{2}(1 + s) ds \leq \frac{3}{4} < 1. \end{aligned}$$

Thus fulfilled each requirement of Theorem 3.1, therefore the validity of solution of the integrodifferential equations are existence and uniqueness.

References

- [1] Altun, D. Turkoglu, B.E. Rhoades, Fixed points of weakly compatible maps satisfying a general contractive condition of integral type, *Fixed Point Theory Appl.*, Vol. 2007, Article ID 17301, 9 pages; doi:10/1155/2007/17301.
- [2] Benmezai, Krasnoselskii-type fixed point theorem in ordered Banach spaces and application to integral equations, *Advances in Pure and Applied Mathematics*, **13**(1)(2022), 50 – 61.
- [3] Branciari, A fixed point theorem for mappings satisfying a general contractive condition of integral type, *Int. J. Math. Math. Sci.*, **29**(9)(2002), 531 – 536.
- [4] B.G. Pachpatte, On a nonlinear Volterra-Fredholm integral equation, *Sarajevo Journal of Mathematics*, **4**(16)(2008), 61 – 71.
- [5] B.G. Pachpatte, On Fredholm type integrodifferential equation, *Tamkang Journal of Mathematics*, **39**(1) (2008), 85.
- [6] B.S. Choudhury, Unique fixed point theorem for weak C-contractive mappings, *Kathmandu Univ. J. Eng. Tech.*, **5**(1)(2009), 6 – 13.
- [7] D. Ilic, V. Rakocevic, Common fixed points for maps on cone metric space, *Journal of Mathematical Analysis and Applications*, **341**(2)(2008), 876 – 882.
- [8] D.P. Shukla, S.K. Tiwari, Unique fixed point theorem for weakly S-contractive mappings, *Gen. Math. Notes*, **4**(1)(2011), 28 – 34.
- [9] D.P. Shukla, S.K. Tiwari, S.K. Shukla, Fixed point theorems for a pair of compatible mappings in integral type equation, *Int. J. of Math. Sci. & Engg. Appls.*, **7**(VI)(2013), 413 – 419.
- [10] G. C. Done, K. L. Bondar, P. U. Chopade, Existence and uniqueness of solution of summation-difference equation of finite delay in cone metric space, *Communications in Mathematics and Applications* **11**(3)(2020), 325 – 334.
- [11] J. Fernandez, N. Malviya, Z.D. Mitrovic, A. Hussain, V. Parvaneh, Some fixed point results on N_b -cone metric spaces over Banach algebra, *Advances in Difference Equations* (2020), <https://doi.org/10.1186/s13662-020-02991-5>.
- [12] J. Fernandez, N. Malviya, A. Savic, M. Paunovic, Z.D. Mitrovic, A. Hussain, V. Parvaneh, The extended cone b-metric-like spaces over Banach algebra and some applications, *Σ Mathematics* (2022) <https://doi.org/10.3390/math10010149>.
- [13] L.G. Huang, X. Zhang, Cone metric spaces and fixed point theorems of contractive mappings, *Journal of Mathematical Analysis and Applications*, **332**(2)(2007), 1468 – 1476.

- [14] M. Abbas, G. Jungck, Common fixed point results for noncommuting mappings without continuity in cone metric spaces, *Journal of Mathematical Analysis and Applications*, **341**(1)(2008), 416 – 420.
- [15] M. Frechet, Sur quelques point du calcul fonctionnel, *Rendiconti del circolo Matematico di palermo*, **22**(1906), 1 – 74, doi:10.1007/BF03018603.
- [16] P. Vijayaraju, B.E. Rhoades, R. Mohanraj, A fixed point theorem for a pair of maps satisfying a general contractive condition of integral type, *Intern. J. Math. Math. Sci.* **15** (2005), 2359-2364.
- [17] P. Raja, S.M. Vaezpour, Some extensions of Banach's contraction principle in complete cone metric spaces, *Fixed Point Theory and Applications*, Volume 2008, Article ID 768294, 11pages.
- [18] T. Abdeljawad, Fixed points for generalized weakly contractive mappings in partial metric spaces, *Math. Comput. Modelling*, **54**(2011), 2923 – 2927.
- [19] S. Aleksic, Z. Kadelburg, Z. D. Mitrovic, S. Radenovic, A new survey: Cone metric spaces, *J. of the Int. Mathematical Virtual Institute*, **(9)**(2019), 93 – 121.
- [20] Z. Kalkan, A. Sahin, Some New Results in Partial Cone b-Metric Space, *Communications in Advanced Mathematical Sciences*, **III**(2)(2020), 67 – 73.